

Precise determination of the branching ratio of the Neutral-Pion Dalitz decay

Husek, Tomáš; Goudzovski, Evgueni; Kampf, Karol

DOI:

[10.1103/PhysRevLett.122.022003](https://doi.org/10.1103/PhysRevLett.122.022003)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Husek, T, Goudzovski, E & Kampf, K 2019, 'Precise determination of the branching ratio of the Neutral-Pion Dalitz decay' Physical Review Letters, vol. 122, no. 2, 022003. <https://doi.org/10.1103/PhysRevLett.122.022003>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Husek et al, Precise determination of the branching ratio of the Neutral-Pion Dalitz decay, Physical Review Letters, Vol. 122, Article No. 022003, Published 18/01/2019, <https://doi.org/10.1103/PhysRevLett.122.022003>

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Precise Determination of the Branching Ratio of the Neutral-Pion Dalitz Decay

Tomáš Husek,^{1,*} Evgueni Goudzovski,^{2,†} and Karol Kampf^{3,‡}

¹IFIC, Universitat de València—CSIC, Apt. Correus 22085, E-46071 València, Spain

²School of Physics and Astronomy, University of Birmingham, Edgbaston, Birmingham B15 2TT, United Kingdom

³Institute of Particle and Nuclear Physics, Faculty of Mathematics and Physics, Charles University, V Holešovičkách 2, 18000 Praha 8, Czech Republic



(Received 13 September 2018; published 18 January 2019)

We provide a new value for the ratio $R = \Gamma(\pi^0 \rightarrow e^+e^-\gamma)/\Gamma(\pi^0 \rightarrow \gamma\gamma) = 11.978(6) \times 10^{-3}$, which is by 2 orders of magnitude more precise than the current Particle Data Group average. It is obtained using the complete set of the next-to-leading-order radiative corrections in the QED sector, and incorporates up-to-date values of the π^0 -transition-form-factor slope. The ratio R translates into the branching ratios of the two main π^0 decay modes: $\mathcal{B}(\pi^0 \rightarrow \gamma\gamma) = 98.8131(6)\%$ and $\mathcal{B}(\pi^0 \rightarrow e^+e^-\gamma) = 1.1836(6)\%$.

DOI: 10.1103/PhysRevLett.122.022003

Introduction.—The present 3% experimental precision on $\mathcal{B}(\pi_D^0)$ [1] (We use the shorthand notation $\pi_f^0 := \pi^0 \rightarrow f$, with $D := e^+e^-\gamma$ and $DD := e^+e^-e^+e^-$.) represents a limitation for rare- π^0 -decay measurements, which commonly use the Dalitz decay π_D^0 for normalization, and is also becoming a limiting factor for rare-kaon-decay measurements. An example is the $K^+ \rightarrow \pi^+e^+e^-$ decay [2]: accurate knowledge of $\mathcal{B}(\pi_D^0)$ would improve the precision on the rate measurement by 30%, and the precision on the low-energy parameter a_+ [3] by 10%. The uncertainty on $\mathcal{B}(\pi_D^0)$ also dominates the precision on the $K^\pm \rightarrow \pi^\pm\pi^0e^+e^-$ rate measurement [4], and is among the principal contributions to the uncertainties on the measured $K_{L,S} \rightarrow \pi^+\pi^-e^+e^-$ rates [5]. In these circumstances, considering the improving precision on rare-decay measurements, and the recent progress on the π^0 -form-factor measurement [6] and radiative corrections for the π_D^0 decay [7], both a precision measurement of $\mathcal{B}(\pi_D^0)$ and an updated theoretical evaluation of this quantity are becoming more important.

Branching ratios can serve to translate lifetimes into decay widths and vice versa. There are several methods to determine the π^0 lifetime: a direct average-distance measurement of the highly relativistic pion, the conserved-vector-current hypothesis connecting the vector form factor (i.e., charged pions) to the π^0 lifetime [8], and the Primakoff effect [9]. Since 2012 its Particle Data Group (PDG) value settled to $\tau_{\pi^0}^{\text{PDG}} = 8.52(18) \times 10^{-17}$ s [10]. Presently, the most precise

π^0 -lifetime measurements are given by two different methods: $\tau_{\pi^0}^{\text{PrimEx}} = 8.32(23) \times 10^{-17}$ s [11] (Primakoff effect, PrimEx experiment at JLab) and $\tau_{\pi^0}^{\text{CERN}} = 8.97(28) \times 10^{-17}$ s [12] (direct measurement, CERN). It is clear that the situation is unsatisfactory and a new independent measurement is desirable. For the Primakoff-effect-type π^0 -lifetime measurements, $\mathcal{B}(\pi_{2\gamma}^0)$ constitutes an essential input.

In this work we discuss the theoretical determination of the following ratio of decay widths:

$$R \equiv \frac{\Gamma(\pi^0 \rightarrow e^+e^-\gamma)}{\Gamma(\pi^0 \rightarrow \gamma\gamma)} = \frac{\mathcal{B}(\pi_D^0)}{\mathcal{B}(\pi_{2\gamma}^0)}. \quad (1)$$

The current PDG value $R|_{\text{PDG}} = 11.88(35) \times 10^{-3}$ is an average of experimental results, the most recent of which comes from 2008 and is based on archived ALEPH data [13]. Other measurements with competitive uncertainties date back to 1981 [14] and 1961 [15]. The branching ratios $\mathcal{B}(\pi_{2\gamma}^0)|_{\text{PDG}} = 98.823(34)\%$ and $\mathcal{B}(\pi_D^0)|_{\text{PDG}} = 1.174(35)\%$ [1] are subsequently obtained from a constrained fit.

Besides the direct extraction of R from experiment, the shape of the singly virtual π^0 transition form factor (TFF) can be measured. This can be expanded in the transferred momentum squared, with the linear coefficient called the (TFF) slope a_π . Since the slope embodies the most relevant input to the ratio R regarding the (nonperturbative) low-energy QCD sector [the peculiar [16] TFF normalization $\mathcal{F}(0)$ conveniently drops out], its knowledge from experiment is crucial to obtain a model-independent prediction of R . Recently, it was measured in the NA62 experiment, which analyzed 1.1×10^6 fully reconstructed π_D^0 decays with the result $a_\pi^{\text{NA62}} = 3.68(57)\%$ [6], taking into account the complete set of next-to-leading-order (NLO) radiative corrections in the QED sector [7]. The current PDG value is

Published by the American Physical Society under the terms of the Creative Commons Attribution 4.0 International license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.

dominated by two inputs: the above NA62 result and the value provided by the CELLO Collaboration [$a_\pi^{\text{CELLO}} = 3.26(37)\%$] [17] by (model-dependent) extrapolation from the spacelike region.

Our calculation combines a wide range of theoretical models and available experimental results on the TFF shape and the well-established QED calculation including the complete set of NLO corrections, taking into account higher orders in the QED expansion in a conservative uncertainty estimate. As such, it represents a precise and reliable improvement (by two orders of magnitude) to the current PDG-based value of R , which might be further used in various theoretical predictions and experimental analyses. Moreover, for the first time, the slope corrections were not neglected in the bremsstrahlung contribution. Finally, we present R for the full as well as partial kinematic regions.

Measurements of the TFF shape or the ratio R require significant theoretical input and depend crucially on the proper incorporation of radiative corrections. Consequently, a statement that experiment itself provides a more relevant value of R than our theoretical prediction is by its nature imprecise. However, the computation would not be possible without the experimental evidence that the TFF slope lies within a certain range of values. An example of how theoretical inputs influence the experimental values in this sector is the well-known discrepancy in the rare decay $\pi_{e^+e^-}^0$ driven most probably by the approximate radiative corrections [18] which do not agree with the exact calculation [19]; for details and discussion see Refs. [20,21].

Theoretical framework.—Considering the QED expansion, the leading-order (LO) π^0 -Dalitz-decay differential width reads [7,22,23]

$$\frac{d^2\Gamma^{\text{LO}}(x, y)}{dx dy} = \frac{\alpha}{\pi} \Gamma(\pi^0 \rightarrow \gamma\gamma) \left| \frac{\mathcal{F}(M_\pi^2 x)}{\mathcal{F}(0)} \right|^2 \frac{(1-x)^3}{4x} \left(1 + y^2 + \frac{4m_e^2}{M_\pi^2 x} \right), \quad (2)$$

where the two-photon decay width is parametrized as

$$\Gamma(\pi^0 \rightarrow \gamma\gamma) \equiv \frac{e^4 M_\pi^3}{64\pi} |\mathcal{F}(0)|^2. \quad (3)$$

TABLE I. The slope and curvature of the singly virtual pion TFF in various approaches. The values given by the VMD, LMD, and THS models are compared with the results of the recent dispersive calculation [36,37] incorporating inputs from both the space- and timelike regions (and updating Ref. [38]), the method of Padé approximants [39] mainly based on the extrapolation of the spacelike data (as was the previous work [40]) and supported by the low-energy timelike data, the recent measurement performed by the NA62 experiment [6], or the PDG average [1]. Inherent model uncertainties (due to large- N_c and chiral limits) are not fully included in the THS value [21]. Additionally, a recent time-like-region measurement by the A2 Collaboration reads $a_\pi^{\text{A2}} = 3.0(1.0)\%$ [41].

Source	VMD	LMD	THS	Dispers.	Padé aps.	NA62	PDG
$a_\pi[\%]$	3.00	2.45	2.92(4)	3.15(9)	3.21(19)	3.68(57)	3.35(31)
$b_\pi[10^{-3}]$	0.90	0.74	0.87(2)	1.14(4)	1.04(22)

Above, M_π and m_e are the neutral-pion and electron masses, respectively. The definition (3) holds to all orders in the QED and chiral perturbation theory expansions [16] and covers also possible physics from beyond the standard model, simply putting these nontrivial dynamical effects into the TFF normalization $\mathcal{F}(0)$. As usual, kinematical variables x and y are defined as

$$x = \frac{(p_{e^-} + p_{e^+})^2}{M_\pi^2}, \quad y = -\frac{2}{M_\pi^2} \frac{p_{\pi^0} \cdot (p_{e^-} - p_{e^+})}{(1-x)}, \quad (4)$$

with p denoting four-momenta of respective particles.

The slope a_π and curvature b_π of the singly virtual pion TFF are defined in terms of the Taylor expansion in the invariant mass of the vector current [24,25]:

$$\left| \frac{\mathcal{F}(M_\pi^2 x)}{\mathcal{F}(0)} \right| \equiv f(x) = 1 + a_\pi x + b_\pi x^2 + \mathcal{O}(x^3). \quad (5)$$

This parametrization is sufficient in the whole (small) region relevant to the π_D^0 decay. Having particular theoretical models at hand one can immediately explore the properties of $\mathcal{F}(q^2)$ and calculate a_π and b_π . As examples we briefly mention the vector-meson-dominance (VMD) ansatz [25,26] together with the lowest-meson-dominance (LMD) [27,28] and two-hadron-saturation (THS) [21] models; see Ref. [29] for more details. These belong to a family of large- N_c motivated analytic resonance-saturation models and as such they can be straightforwardly used in the calculation of radiative corrections. By means of the first and second derivatives it is easy to find the analytic expressions for a_π and b_π within these models; for details see Sec. 5 of Ref. [21]. Numerical results are shown in Table I, together with other theoretical approaches and experimental results.

From Eq. (5) it follows that

$$f^2(x) = 1 + 2a_\pi x + (a_\pi^2 + 2b_\pi)x^2 + \mathcal{O}(x^3). \quad (6)$$

We can use the expansion (6) to obtain a simple formula for the LO width. Inserting Eq. (6) into Eq. (2) and taking into account that $x \in (4m_e^2/M_\pi^2, 1)$ and $y \in (-\beta(M_\pi^2 x), \beta(M_\pi^2 x))$ with $\beta(s) \equiv \sqrt{1 - 4m_e^2/s}$, we get

$$R^{\text{LO}} = \frac{\alpha}{\pi} \left[\frac{4}{3} \ln \frac{M_\pi}{m_e} - \frac{1}{3} (7 - a_\pi) + \frac{1}{30} (a_\pi^2 + 2b_\pi) + (12 - 8a_\pi) \frac{m_e^2}{M_\pi^2} + \mathcal{O}\left(\frac{m_e^4}{M_\pi^4}\right) \right]. \quad (7)$$

This expression is a very good approximation with the precision $\eta(R^{\text{LO}}) \simeq 10^{-9}$ (evaluated for parameters close to the physical ones). As is a common practice, e.g., in Ref. [6], further on the prescription $f^2(x)|_{\text{lin}} \equiv (1 + a_\pi x)^2$ is used, which is accurate to first order in x . The effect of the missing $2b_\pi x^2$ term is negligible, since the region where such a difference arises (i.e., when $x \simeq 1$) is suppressed; cf. Eq. (2) and also Table III later on. If we consider $b_\pi \simeq a_\pi^2$, as suggested by the models (cf. Table I), we introduce an error of $\sigma(a_\pi) \simeq a_\pi^2/5$, i.e., a relative error $\eta(a_\pi) \lesssim 1\%$, on the estimate of a_π being well under the current experimental precision. The previous discussion also implies that the effect of the $a_\pi x$ term on the Dalitz-decay rate is limited, letting us provide a very precise determination of R while allowing for 20% uncertainty on a_π . Finally, let us note that dropping b_π out of Eq. (7) decreases its precision down to $\eta(R^{\text{LO}}|_{b_\pi=0}) \simeq 10^{-5}$, being still a good approximation for our purpose in view of the above discussion.

In the rest of the section we address the NLO QED-sector radiative corrections; see Fig. 1 for Feynman diagrams. It is convenient to introduce the NLO correction δ to the LO differential width [and thus to write schematically $d\Gamma = (1 + \delta + \dots)d\Gamma^{\text{LO}}$], which can be in general (for the two- and onefold differential case, respectively) defined as

$$\delta(x, y) = \frac{d^2\Gamma^{\text{NLO}}}{dx dy} \bigg/ \frac{d^2\Gamma^{\text{LO}}}{dx dy}, \quad \delta(x) = \frac{d\Gamma^{\text{NLO}}}{dx} \bigg/ \frac{d\Gamma^{\text{LO}}}{dx}. \quad (8)$$

One can obtain $\delta(x)$ from $\delta(x, y)$ using the following prescription:




FIG. 1. NLO QED radiative corrections for π^0 : (a) vacuum-polarization insertion; (b),(c) one-loop $1\gamma\text{IR}$ contribution; (d) vertex correction; (e) bremsstrahlung.

$$\delta(x) = \frac{3}{8} \frac{1}{\beta(M_\pi^2 x)} \left(1 + \frac{2m_e^2}{M_\pi^2 x} \right)^{-1} \times \int_{-\beta(M_\pi^2 x)}^{\beta(M_\pi^2 x)} \delta(x, y) \left(1 + y^2 + \frac{4m_e^2}{M_\pi^2 x} \right) dy. \quad (9)$$

To calculate the NLO radiative corrections we use the approach documented in Refs. [7,42], which reviewed and extended the classical work of Mikaelian and Smith [22]. Hence, together with the bremsstrahlung (BS) beyond the soft-photon approximation, we take into account in the following calculations the one-photon-irreducible ($1\gamma\text{IR}$) contribution; see Figs. 1(b) and 1(c).

For historical reasons [22,23,43–45], let us discuss the case when the $1\gamma\text{IR}$ contribution to the NLO radiative corrections is not considered in the analysis to extract the TFF slope from the data. If we start with the equation among the measured spectral shapes (onefold differential widths) and eliminate $d\Gamma^{\text{LO}}/dx|_{f(x)=1}$ from both sides, take the expansion (6) to the linear order, and neglect the corrections of order αa_π , we find

$$\Delta a_\pi x \equiv (a_\pi - a_\pi^{1\gamma\text{IR}})x \simeq -\frac{1}{2} \delta_{1\gamma\text{IR}}^{\text{NLO}}(x), \quad x \ll 1. \quad (10)$$

Numerically, $\Delta a_\pi \simeq 0.5\%$ [23]. This is the value to be added to the experimental value $a_\pi^{1\gamma\text{IR}}$ (extracted neglecting the $1\gamma\text{IR}$ contribution) in order to find an estimate of the pure-low-energy-QCD-sector parameter a_π with all the QED radiative corrections subtracted. Above, $\delta_{1\gamma\text{IR}}^{\text{NLO}}(x)$ is calculated from $\delta^{1\gamma\text{IR}}(x, y)$ stated in Sec. IV of Ref. [7] using the prescription (9). Note that the $1\gamma\text{IR}$ contribution was already taken into account in the NA62 analysis [6] and a_π^{NA62} does not need to be corrected by Δa_π .

Finally, taking the prescription $f^2(x)|_{\text{lin}}$ and the NLO QED radiative corrections to approximate the exact Dalitz-decay differential width beyond LO (and consequently $\Delta R \equiv R - R^{\text{LO}}$) we arrive at

$$\Delta R \simeq R^{\text{NLO}} \equiv \frac{\alpha}{\pi} \iint (1 + a_\pi x)^2 \delta(x, y) \times \frac{(1-x)^3}{4x} \left(1 + y^2 + \frac{4m_e^2}{M_\pi^2 x} \right) dx dy. \quad (11)$$

Calculation and uncertainty estimation.—Our aim now is to precisely and reliably (using conservative error estimates) determine R . In the following we choose $a_\pi^{\text{univ}} \equiv 3.55(70)\%$ by stretching the uncertainty band over the whole interval of values suggested by different approaches; cf. Table I. From Eq. (7) we get

$$R^{\text{LO}} = 11.879(5) \times 10^{-3}, \quad (12)$$

and based on Eq. (11) we arrive at

TABLE II. Individual contributions of the NLO radiative corrections for R in $[10^{-3}]$. The virt label stands for virtual corrections [Figs. 1(a) and 1(d)] and div (conv) for the divergent (convergent) parts of the bremsstrahlung contribution [Fig. 1(e)] [46]. The listed uncertainties stem from the uncertainty of a_π^{univ} . In the case of the $1\gamma\text{IR}$ correction [Figs. 1(b) and 1(c)], a particular model for the doubly virtual TFF (LMD, etc.) is necessary to introduce. The resulting model dependence is suppressed [7] and related uncertainty included.

$R_{\text{virt}}^{\text{NLO}}$	$R_{\text{BS,conv}}^{\text{NLO}}$	$R_{\text{BS,div}}^{\text{NLO}}$	$R_{1\gamma\text{IR}}^{\text{NLO}}$	R^{NLO}
-0.0750(2)	-0.15759(2)	0.3363(3)	-0.00466(2)	0.09911(7)

$$\Delta R = 0.099(3) \times 10^{-3}. \quad (13)$$

During the estimation of the above uncertainty, higher orders in the QED expansion were considered, surpassing in size the uncertainty stemming from the TFF dependence. In this regard, we can take the absolute value of the *dominant* correction ($R_{\text{BS,div}}^{\text{NLO}}$; see Table II) as the typical expected maximum value appearing in NLO and anticipate that the NNLO correction is suppressed compared to the NLO one in the similar manner as NLO is suppressed with respect to LO: circa on the level of 3%. This uncertainty is already conservative: the total NLO correction is on the level of 1%. Summing Eqs. (12) and (13) we finally obtain

$$R = 11.978(5)(3) \times 10^{-3}. \quad (14)$$

This is one of the main results of the presented work. The former uncertainty stands for the TFF effects and the latter for neglecting the higher-order corrections; [Relaxing the requirement of providing a conservative value, one can significantly reduce the former uncertainty (stemming from the TFF effects) taking into account a particular result from Table I: e.g., with the most precise entry—the dispersion-theoretical result [36,37]—by a factor of 8. Higher-order QED corrections would need to be computed to achieve an additional gain of precision.] m_e , M_π , and α are known very precisely.

TABLE III. The values of $R(x_{\text{cut}})$ for chosen sample values of x_{cut} . To be suitable for interpolation, higher precision is used. The quoted uncertainties are dominated by the TFF-slope knowledge (for its value we assume a_π^{univ}); the additional 3% uncertainty covering the higher-order corrections is also included. Note different additional multiplicative factors depending on x_{cut} .

x_{cut}	$R(x_{\text{cut}})[10^{-5}]$	x_{cut}	$R(x_{\text{cut}})[10^{-5}]$	x_{cut}	$R(x_{\text{cut}})[10^{-6}]$	x_{cut}	$R(x_{\text{cut}})[10^{-8}]$
0.05	203.72(45)	0.30	22.13(13)	0.55	24.20(22)	0.80	67.28(85)
0.10	117.03(36)	0.35	14.787(97)	0.60	14.07(14)	0.85	19.81(27)
0.15	74.38(29)	0.40	9.756(71)	0.65	7.703(80)	0.90	3.594(54)
0.20	49.11(23)	0.45	6.313(50)	0.70	3.890(43)	0.95	0.1967(35)
0.25	32.92(17)	0.50	3.978(34)	0.75	1.757(21)	1.00	0

This calculation also includes all contributions from the decays where additional photon(s) with arbitrarily high (kinematically allowed) energies are radiated. Indeed, the bremsstrahlung correction at NLO (calculated *à la* Refs. [7,22]) takes into account an additional final-state photon and integrates over its energy and emission angle without any additional cuts. The results are thus meant to be used for the *inclusive* process. However, quantities for *exclusive* processes can be obtained in a similar way while introducing some specific cutoff in the bremsstrahlung correction $\delta^{\text{BS}}(x, y)$. A combined approach was used in the analysis of the recent NA62 measurement [6], when an additional photon was simulated above the cutoff given by the detector sensitivity. To conclude, for each experimental setup the specific approach for including radiative corrections must be used. When it applies, we explicitly state (as in the abstract) that the results include an additional final-state photon, denoting it as (γ) . We also take this tacitly into account in the results for R , e.g., in Eq. (14).

In experiments, specific kinematic regions might be considered. The sample values for

$$R(x_{\text{cut}}) \equiv \frac{\mathcal{B}(\pi^0 \rightarrow e^+ e^- \gamma(\gamma), x > x_{\text{cut}})}{\mathcal{B}(\pi_{2\gamma}^0)} \quad (15)$$

are listed in Table III, using which one can also obtain values for any intermediate region. As an example, in the π^0 -rare-decay measurement performed by KTeV [47] the region $x > x_{\text{cut}} = 0.232$ was used for the Dalitz decay, which served as the normalization channel in this search. The direct calculation based on this work leads to $R(0.232) = 0.380(2) \times 10^{-3}$ and the interpolation based on Table III gives $R(0.232)|_{\text{intpol}} = 0.379(2) \times 10^{-3}$, which is compatible within uncertainties. In Ref. [48] the value $[R(0.2319)/R]|_{\text{KTeV}} = 0.0319$ was used, which is compatible with our calculation: $R(0.232)/R = 0.0317(2)$.

Based on R , we can predict $\mathcal{B}(\pi_{2\gamma}^0)$ and $\mathcal{B}(\pi_{\text{D}}^0)$. Considering the uncertainty of R , we can write

$$1 - \mathcal{B}(\pi_{\text{DD}}^0) \simeq \mathcal{B}(\pi_{2\gamma}^0) + \mathcal{B}(\pi_{\text{D}(\gamma)}^0), \quad (16)$$

since the branching ratios of other decay modes are smaller than 10^{-6} . Using $\mathcal{B}(\pi_{\text{DD}}^0) = 3.3(2) \times 10^{-5}$ [1,49] (double-Dalitz decay), we find

$$\mathcal{B}(\pi_{2\gamma}^0) \simeq \frac{1 - \mathcal{B}(\pi_{\text{DD}}^0)}{1 + R} = 98.8131(6)\%. \quad (17)$$

Note that taking tacitly into account inclusive Dalitz decays in Eq. (16) is justified and contributes to the relevant decay modes. Finally, the Dalitz-decay branching ratio reads

$$\mathcal{B}(\pi_{\text{D}(\gamma)}^0) \simeq \frac{R}{1 + R} [1 - \mathcal{B}(\pi_{\text{DD}}^0)] = 1.1836(6)\%. \quad (18)$$

The above results are compatible with the PDG averages, exhibiting much higher precision.

Let us see how the new result on the Dalitz-decay branching ratio (18) influences a completely different family of processes on a simple example of the $K^+ \rightarrow \pi^+ e^+ e^-$ decay measurements. The low-energy parameters a_+ and b_+ were measured by the NA48/2 collaboration to be $a_+ = -0.578(16)$ and $b_+ = -0.779(66)$, leading to the model-dependent branching ratio $\mathcal{B}(K^+ \rightarrow \pi^+ e^+ e^-) = 3.11(12) \times 10^{-7}$, using the 2008 PDG average $\mathcal{B}(\pi_{\text{D}}^0) = 1.198(32)\%$ [50] for normalization [2]. The central value of our result (18) is 1.2% lower than the quoted PDG average and has a negligible error. The remaining external uncertainty on the measurement [2] related to the normalization comes from $\mathcal{B}(K^+ \rightarrow \pi^+ \pi^0)$ known to 0.4% precision. The corrected values are $a_+ = -0.575(14)$, $b_+ = -0.771(64)$, and $\mathcal{B}(K^+ \rightarrow \pi^+ e^+ e^-) = 3.07(10) \times 10^{-7}$. Note that considering the external errors on a_+ and b_+ quoted in Ref. [2], further experimental progress on $K^+ \rightarrow \pi^+ e^+ e^-$ measurement would be impossible without improvement on $\mathcal{B}(\pi_{\text{D}}^0)$.

Comparison and conclusion.—Radiative corrections for the integral decay width were first addressed by Joseph [51], who numerically arrived to $\Delta R|_{\text{jph}} = 0.105 \times 10^{-3}$ neglecting, among others, the pion TFF slope. A simple analytical prescription in the limit of the vanishing electron mass was later found by Lautrup and Smith [52]:

$$\Delta R|_{\text{L\&S}} = \left(\frac{\alpha}{\pi}\right)^2 \left[\frac{8}{9} \ln^2 \frac{M_\pi}{m_e} - \frac{1}{9} (19 - 4a_\pi) \ln \frac{M_\pi}{m_e} + 2\zeta(3) - \frac{2}{27} \pi^2 + \frac{137}{81} - \frac{63}{108} a_\pi + \mathcal{O}\left(\frac{m_e}{M_\pi}\right) \right]. \quad (19)$$

Numerically, $\Delta R|_{\text{L\&S}}^{a_\pi=0} = 0.10378 \times 10^{-3}$ and $\Delta R|_{\text{L\&S}}^{a_\pi^{\text{univ}}} = 0.10414(7) \times 10^{-3}$. The two approaches are compatible and should be compared with our result (13). However, the $1\gamma\text{IR}$ contribution was, due to inappropriate assumptions and arguments based on Low's theorem [53–55], considered negligible and left out; see also Refs. [22,44]. The exact calculation shows its significance [7,23,43,45] and it

thus embodies the main source of the difference between our result and the previous works. Moreover, the symmetrization with respect to the two photons in the bremsstrahlung contribution was neglected in Refs. [51,52]. This interference of the diagrams from Fig. 1(e) is indeed negligible and corresponds (for $a_\pi = 0$) to $\Delta R_{\text{interf}}^{\text{BS}} = 0.000360 \times 10^{-3}$. Let us stress again that the prediction (14) is based on the complete calculation which includes the entire bremsstrahlung and $1\gamma\text{IR}$ contributions. Here, TFF effects were taken into account also in the bremsstrahlung correction [46] and the mass of the final-state leptons was *not* neglected.

Our main result (14) together with the value (17) should be considered as an alternative to the current PDG averages which opens the way to a new level of precision for a whole class of other processes, for instance, for the already mentioned kaon decays. Similarly, the current situation, when the precision on $\mathcal{B}(\pi_{\text{D}}^0)|_{\text{PDG}}$ dominates the uncertainty on $\mathcal{B}(\pi_{\text{DD}}^0)$ [56] and is the largest source of uncertainty on $\mathcal{B}(\pi_{e^+e^-}^0)$ [47], is improved.

We thank G. D'Ambrosio, M. Hoferichter, and A. Portelli for initial suggestions, P. Sanchez-Puertas for helpful discussions, and J. Portolés for comments on the manuscript. This work has been supported in part by the Agencia Estatal de Investigación (AEI, ES) and the European Regional Development Fund (ERDF, EU) [Grants No. FPA2014-53631-C2-1-P, No. FPA2017-84445-P, and No. SEV-2014-0398], by Generalitat Valenciana [Grant No. PROMETEO/2017/053], by the Czech Science Foundation grant GAČR 18-17224S and by the ERC starting Grant No. 336581 “KaonLepton.”

*thusek@ific.uv.es

†eg@hep.ph.bham.ac.uk

‡karol.kampf@mff.cuni.cz

- [1] M. Tanabashi *et al.* (Particle Data Group), *Phys. Rev. D* **98**, 030001 (2018).
- [2] J. R. Batley *et al.* (NA48/2 Collaboration), *Phys. Lett. B* **677**, 246 (2009).
- [3] G. D'Ambrosio, G. Ecker, G. Isidori, and J. Portolés, *J. High Energy Phys.* **08** (1998) 004.
- [4] J. R. Batley *et al.* (NA48/2 Collaboration), *Phys. Lett. B* **788**, 552 (2019).
- [5] A. Lai *et al.* (NA48 Collaboration), *Eur. Phys. J. C* **30**, 33 (2003).
- [6] C. Lazzeroni *et al.* (NA62 Collaboration), *Phys. Lett. B* **768**, 38 (2017).
- [7] T. Husek, K. Kampf, and J. Novotný, *Phys. Rev. D* **92**, 054027 (2015).
- [8] M. Bychkov *et al.*, *Phys. Rev. Lett.* **103**, 051802 (2009).
- [9] H. Primakoff, *Phys. Rev.* **81**, 899 (1951).
- [10] J. Beringer *et al.* (Particle Data Group), *Phys. Rev. D* **86**, 010001 (2012).
- [11] I. Larin *et al.* (PrimEx Collaboration), *Phys. Rev. Lett.* **106**, 162303 (2011).

- [12] H. W. Atherton *et al.*, *Phys. Lett.* **158B**, 81 (1985).
- [13] A. Beddall and A. Beddall, *Eur. Phys. J. C* **54**, 365 (2008).
- [14] M. A. Schardt, J. S. Frank, C. M. Hoffman, R. E. Mischke, D. C. Moir, and P. A. Thompson, *Phys. Rev. D* **23**, 639 (1981).
- [15] N. P. Samios, *Phys. Rev.* **121**, 275 (1961).
- [16] K. Kampf and B. Moussallam, *Phys. Rev. D* **79**, 076005 (2009).
- [17] H. J. Behrend *et al.* (CELLO Collaboration), *Z. Phys. C* **49**, 401 (1991).
- [18] L. Bergström, *Z. Phys. C* **20**, 135 (1983).
- [19] P. Vařko and J. Novotný, *J. High Energy Phys.* **10** (2011) 122.
- [20] T. Husek, K. Kampf, and J. Novotný, *Eur. Phys. J. C* **74**, 3010 (2014).
- [21] T. Husek and S. Leupold, *Eur. Phys. J. C* **75**, 586 (2015).
- [22] K. O. Mikaelian and J. Smith, *Phys. Rev. D* **5**, 1763 (1972).
- [23] K. Kampf, M. Knecht, and J. Novotný, *Eur. Phys. J. C* **46**, 191 (2006).
- [24] S. Berman and D. Geffen, *Nuovo Cimento* **18**, 1192 (1960).
- [25] L. Landsberg, *Phys. Rep.* **128**, 301 (1985).
- [26] J. J. Sakurai, *Currents and Mesons* (University of Chicago Press, Chicago, 1969).
- [27] S. Peris, M. Perrottet, and E. de Rafael, *J. High Energy Phys.* **05** (1998) 011.
- [28] M. Knecht, S. Peris, M. Perrottet, and E. de Rafael, *Phys. Rev. Lett.* **83**, 5230 (1999).
- [29] The simplest models do not, for instance, satisfy some of the major high-energy constraints: VMD violates operator-product-expansion (OPE) constraints [21,30] and the LMD model cannot satisfy the Brodsky-Lepage scaling limit [31,32]. Although the THS model by construction satisfies the mentioned constraints [21] and surpasses in this sense the VMD and LMD models, it inevitably suffers from shortcomings common to the whole family: for instance, a presence of systematic uncertainties at low energies [33] (see also Ref. [34]) or, at in the high-energy region, $\lim_{q^2 \rightarrow \infty} q^2 \mathcal{F}^{\text{THS}}(p^2, q^2)$ not being finite for all p^2 , as a direct consequence of satisfying OPE [35].
- [30] M. Knecht and A. Nyffeler, *Eur. Phys. J. C* **21**, 659 (2001).
- [31] G. P. Lepage and S. J. Brodsky, *Phys. Lett.* **87B**, 359 (1979).
- [32] G. P. Lepage and S. J. Brodsky, *Phys. Rev. D* **22**, 2157 (1980).
- [33] M. Golterman and S. Peris, *Phys. Rev. D* **74**, 096002 (2006).
- [34] P. Masjuan and S. Peris, *J. High Energy Phys.* **05** (2007) 040.
- [35] S. Leupold and B. Kubis (personal communication).
- [36] M. Hoferichter, B.-L. Hoid, B. Kubis, S. Leupold, and S. P. Schneider, *Phys. Rev. Lett.* **121**, 112002 (2018).
- [37] M. Hoferichter, B.-L. Hoid, B. Kubis, S. Leupold, and S. P. Schneider, *J. High Energy Phys.* **10** (2018) 141.
- [38] M. Hoferichter, B. Kubis, S. Leupold, F. Niecknig, and S. P. Schneider, *Eur. Phys. J. C* **74**, 3180 (2014).
- [39] P. Masjuan and P. Sanchez-Puertas, *Phys. Rev. D* **95**, 054026 (2017).
- [40] P. Masjuan, *Phys. Rev. D* **86**, 094021 (2012).
- [41] P. Adlarson *et al.* (A2 Collaboration), *Phys. Rev. C* **95**, 025202 (2017).
- [42] T. Husek, K. Kampf, J. Novotný, and S. Leupold, *Phys. Rev. D* **97**, 096013 (2018).
- [43] G. B. Tupper, T. R. Grose, and M. A. Samuel, *Phys. Rev. D* **28**, 2905 (1983).
- [44] M. Lambin and J. Pestieau, *Phys. Rev. D* **31**, 211 (1985).
- [45] G. Tupper, *Phys. Rev. D* **35**, 1726 (1987).
- [46] The value corresponding to the correction $\delta_{\text{conv}}^{\text{BS}}$ includes not only the part with neglected slope $\Delta R_{\text{conv}, a_\pi^0=0}^{\text{BS}} = -0.1582(1) \times 10^{-3}$, but newly the form-factor correction was taken into account: $\Delta R_{\text{slope}}^{\text{BS}} = 0.00062(13) \times 10^{-3}$. This was calculated based on the final part of Section V of [7] and is (due to the smallness of a_π) compatible with the approach described in Section V of [42].
- [47] E. Abouzaid *et al.* (KTeV Collaboration), *Phys. Rev. D* **75**, 012004 (2007).
- [48] R. Niclasen, Ph. D. thesis, Colorado University, 2006.
- [49] K. Kampf, J. Novotný, and P. Sanchez-Puertas, *Phys. Rev. D* **97**, 056010 (2018).
- [50] C. Amsler *et al.* (Particle Data Group), *Phys. Lett. B* **667**, 1 (2008).
- [51] D. Joseph, *Nuovo Cimento* **16**, 997 (1960).
- [52] B. Lautrup and J. Smith, *Phys. Rev. D* **3**, 1122 (1971).
- [53] F. E. Low, *Phys. Rev.* **110**, 974 (1958).
- [54] S. L. Adler and Y. Dothan, *Phys. Rev.* **151**, 1267 (1966).
- [55] J. Pestleau, *Phys. Rev.* **160**, 1555 (1967).
- [56] E. Abouzaid *et al.* (KTeV Collaboration), *Phys. Rev. Lett.* **100**, 182001 (2008).